

AD-A211 142 DOCUMENTATION PAGE

Form Approved
OMB No. 3704-0188

2

1a. SECURITY CLASSIFICATION AUTHORITY		1b. RESTRICTIVE MARKINGS DTIC FILE COPY	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAFSAM-PROC-87-16		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION USAF School of Aerospace Medicine	6b. OFFICE SYMBOL (if applicable) USAFSAM/RZP	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Human Systems Division (AFSC) Brooks Air Force Base, TX 78235-5301		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION USAF School of Aerospace Medicine	8b. OFFICE SYMBOL (if applicable) USAFSAM/RZP	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Human Systems Division (AFSC) Brooks Air Force Base, TX 78235-5301		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2312
		TASK NO. W1	WORK UNIT ACCESSION NO. 15
11. TITLE (Include Security Classification) Gel State Chemiluminescence: An Artificial Electron Transport System			
12. PERSONAL AUTHOR(S) Kiel, Johnathan L.; Simmons, David M.; Erwin, David N.			
13a. TYPE OF REPORT Interim	13b. TIME COVERED FROM 86/4/1 TO 87/4/1	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
06	07		
20	14		
		/Gel; Chemiluminescence Peroxidase; Oxidase	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>Crosslinked glucose oxidase and horseradish peroxidase can mimic the macrophage oxidative function in vivo in respect to tumor cells as targets. We used crosslinked glucose oxidase, horseradish peroxidase, and luminol-bovine serum albumin entrapped in filter paper disks to study the chemiluminescent properties of this macrophage-mimicking material. The disks showed a sensitivity to D-glucose down to 1 ug/ml. The luminescence was driven specifically by glucose at pH 6.0, 6.5, 7.0, 7.4, 8.0, and 10.0. The best signal-to-noise ratio, 44,705 to 1, was achieved at pH 7.4. Superoxide dismutase enhanced the chemiluminescence and catalase suppressed it. Therefore, superoxide was produced by the crosslinked system, and the chemiluminescence was dependent on hydrogen peroxide.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Johnathan L. Kiel		22b. TELEPHONE (Include Area Code) (512) 536-3583	22c. OFFICE SYMBOL USAFSAM/RZP

From: OXYGEN RADICALS IN BIOLOGY AND MEDICINE
Edited by Michael G. Simic, Karen A. Taylor,
John F. Ward, and Clemens von Sonntag
(Plenum Publishing Corporation, 1988)

GEL STATE CHEMILUMINESCENCE: AN ARTIFICIAL ELECTRON TRANSPORT SYSTEM*

Johnathan L. Kiel, David M. Simmons, and David N. Erwin

Radiation Sciences Division (RZP)
USAF School of Aerospace Medicine
Brooks AFB, Texas 78235-5301, U.S.A.

INTRODUCTION

Crosslinked glucose oxidase (GO) and horseradish peroxidase (HRP) can mimic macrophage oxidative function in vivo in respect to tumor cells as targets.¹⁻³ Both the flavoprotein of NADPH oxidase of macrophages and GO contain flavin adenine dinucleotide as their prosthetic group.⁴ HRP has a noncovalently bound hematoporphyrin IX prosthetic group as does the cytochrome b₅₅₉ (green hemoprotein) of NADPH oxidase.⁴ Therefore, we anticipated a similarity in biochemical mechanism of action and chemiluminescence (CL) between the semi-synthetic crosslinked GO and HRP and the macrophage NADPH oxidase.

RESULTS AND DISCUSSION

We crosslinked 25 μ l of GO (1 mg/ml), HRP (1 mg/ml), and luminol (LH₂)-bovine serum albumin (30 mg/ml) with 0.5% glutaraldehyde in pH 6.9 PBS in a 7-mm diameter filter paper disk to study the CL properties of this macrophage mimicking material (IC-1L).^{1,2,5}

Table 1 shows that the CL (recorded with a Turner Designs 20e luminometer) of the disks was driven specifically by glucose (200 μ l of 0.5 mg/ml) at pH 6.0, 6.5, 7.0, 7.4, 8.0, and 10.0. The best signal-to-noise ratio, 44,705 to 1, was achieved at pH 7.4. Even without luminol, there was considerable CL from IC-1 (189 to 1 signal-to-noise ratio) when disks treated with glucose were compared with those without glucose at pH 7.4.

The kinetics of the CL also varied with age and handling of the disks. Disks produced maximum CL on the second day after preparation, as did those in Table 1. The peak CL was 6 minutes after activation and persisted for about 3 minutes. On the first day, third day, and subsequent days after synthesis and storage at 4°C, the peak CL of the disks was replaced by a plateau at a lower maximum CL that was achieved slowly

*By acceptance of this article, the publisher or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering the article.

Table 1. Chemiluminescence (relative light units/45 secs \pm S.D. after 6 min) of Crosslinked Glucose Oxidase, Horseradish Peroxidase, and Bovine Serum Albumin (IC-1) or Luminol-Bovine Serum Albumin (IC-1L)

pH ^a	IC-1	IC-1L
6.0	0.365 \pm 0.0146(n=5)	6.851 \pm 0.862(n=6)
6.5	0.370 \pm 0.0204(n=5)	16.76 \pm 1.778(n=7)
7.0	1.283 \pm 0.102(n=5)	112.4 \pm 1.329(n=6)
7.4	1.477 \pm 0.181(n=4)	348.7 \pm 11.57(n=7)
7.4(NG) ^b	---	0.0078 \pm 0.0055(n=5)
8.0	1.773 \pm 0.337(n=4)	1998.0 \pm 228.5(n=6)
8.0(NG)	---	0.4865 \pm 0.008(n=6)
10.0	1.045 \pm 0.068(n=4)	2397.0 \pm 197.0(n=7)
10.0(NG)	---	0.8265 \pm 0.011(n=6)

^aThe buffer was phosphate buffered saline (0.01 M sodium phosphate and 0.154 M NaCl) with 0.5 mg/ml D-glucose; at pH 8.0 and 10.0 the PBS pH was adjusted with 0.01 M sodium carbonate.

^bNG = PBS without glucose.

and persisted for 6 to 10 minutes (Figure 1). Also, the spontaneous CL of IC-1L increased with time as the disk remained at room temperature in pH 7.4 PBS. Luminometry with the Quantitative Luminescence Imaging System (QLIS, USAF) indicated that intense localized long-term (30 minutes or more) CL was produced by IC-1L when compared with standard (1×10^{-7} M to 1×10^{-6} M) luminol in dimethyl sulfoxide solutions activated by 0.1 N NaOH (1-2 second CL lifetimes).

In Table 2, we report that the increase in bulk buffer pH around the IC-1L gel (a change of 2.6 units) led to an increase of only 1.68 pH




Figure 1. Chemiluminescence of IC-1L disks at room temperature. Each disk was either activated with 200 µl of pH 7.4 PBS containing 100 µg D-glucose or treated with PBS without glucose.

Table 2. Buffering Capacity and Proton Exchange by IC-1L Disks

Components ^a	pH \pm S.D.	
Buffer (no gel)	7.4	10.0
Bulk solution		
with gel	7.61 \pm 0.04(n=4)	9.09 \pm 0.12(n=3)
with gel and glucose	7.66 \pm 0.05(n=4)	9.12 \pm 0.12(n=3)
Gel surface		
no glucose	6.11 \pm 0.07(n=8)	6.98 \pm 0.02(n=3)
glucose	5.42 \pm 0.09(n=4)	6.90 \pm 0.03(n=3)

^aEach 7-mm paper disk containing 25 μ l of gel was treated with 200 μ l buffer (0.01 M PBS at pH 7.4 or 0.01 M TRIS/saline at pH 10.0) with or without 0.5 mg/ml D-glucose.

units in the surface of activated gel and 1.76 units in gel without glucose. The gel acted like a buffer with a pKa near 5 or 6 as indicated in Table 2. The addition of glucose consistently decreased the surface pH in either pH 7.4 or pH 10.0 buffer. These data indicate that the polyanionic gel entraps protons (hydrogen ions) even though the bulk solution may be very basic. Activation of CL facilitates the accumulation of protons in the gel.

Figure 2 illustrates the nonspecific and specific antioxidative response of IC-1L to inert human serum albumin (HSA), superoxide dismutase (SOD), and catalase (CAT). Comparing Figures 1 and 2, one can see that at 2 μ g, HSA does not affect CL. However, at 2 μ g, SOD enhances the CL. At 20 μ g, HSA quenches CL by 81%, but CAT suppresses CL by 96%. These data indicate a strong participation of H_2O_2 and a minor but significant abortive role of free $\cdot O_2^-$. Dismutation of released $\cdot O_2^-$ facilitates the CL.






Figure 2. Chemiluminescence of IC-1L disks activated as described in Figure 1, but with human serum albumin, superoxide dismutase, or catalase.

CONCLUSIONS

Based on the data we present here, we conclude that the CL mechanism of IC-1L is as follows (where LO_2 is peroxylyminol or cyclooxyluminol and AP^{-2} is aminophthalate):



ACKNOWLEDGEMENTS

This work was sponsored in part by the U.S. Air Force Office of Scientific Research.

REFERENCES

1. J.L. Kiel, " Insoluble Crosslinked Cytotoxic Oxidase-Peroxidase System," U.S. Patent 4,486,408. Dec 4, 1984 (filed Apr 7, 1981).
2. K.E. Everse, H. Lin, E.L. Stuyt, J.C. Brady, F. Buddingh, and J. Everse, Antitumor activity of peroxidase, Brit. J. Cancer 51:743 (1985).
3. C.F. Nathan and Z.A. Cohn, Antitumor effects of hydrogen peroxide in vivo, J. Exp. Med. 154:1539 (1981).
4. R. Lutter, M.L.J. van Schaik, R. van Zwieten, R. Wever, D. Roos, and M.N. Hamers, Purification and partial characterization of the b-type cytochrome from human polymorphonuclear leukocytes, J. Biol. Chem. 260:2237 (1985).
5. J.L. Kiel, Microwave effects on immobilized peroxidase chemiluminescence, Bioelectromag. 4:193 (1983).



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1 20	